

Piperazine 10% syrup

Composition:

Each ml syrup contains:
100 mg piperazine (as citrate)

Indications:

Piperazine is a safe and very effective anthelmintic, active against *Ascaris* infections, as well as *Oxyuris* and *Trichonema* in horses, nodular worms (*Oesophagostomum*) in pigs and ruminants, and hookworms in dogs and cats. It has a partial effect on *Strongylus vulgaris* in horses, *Nematodirus* and *Cooperia* spp in ruminants. No effects are found on flukes or tapeworms. In *Ascaris* infections, piperazine is effective against all stages of the worms. It paralyzes the worms in the intestines, which are then expelled by peristalsis. Migrating larvae are not affected. Therefore a second dose after one week is recommended. In case of lung infections, migrating larvae are however eliminated right from the first dose.

Piperazine syrup is very safe and almost non-toxic, even in very high dosage. It is indicated against *Ascaris* and *Capillaria* in poultry and pigeons, strongylosis and oxyurosis in rabbits and horses, *Ascaris* in horses, *Ascaris*, *Nematodirus*, *Ostertagia*, *Cooperia* and *Oesophagostomum* in cattle, *Oesophagostomum* and *Ascaris* in pigs and hookworms in dogs and cats.

Administration/dosage:

Poultry: 2 ml per liter of drinking water for 1-2 days.

Dogs and cats: 1-2 ml/kg b.w.

Other animals: 100-300 mg of piperazine citrate (1-3 ml) per kg b.w., to be repeated after 3 weeks.

Advised withdrawal times:

meat : 2 days

eggs : 2 days

Conservation:

Dry, cool (8-15°C), protected from light.

Presentation:

bottles 100, 500 ml,

cans of 1000, 5000 ml.