

Oxytetracycline 10%

Composition:

Each ml solution contains:

100 mg oxytetracycline HCl in propylene-glycol carrier

Action:

Broad spectrum antibiotic with bacteriostatic action against Gram-positive and Gram-negative germs such as E. coli, Pasteurella, Salmonella, Brucella, Streptococcus, Clostridium, Haemophilus, Corynebacterium, Anthrax, Staphylococcus. Rickettsia, Mycoplasma, Spirochaetes, and Actinomyces are affected. After parenteral administration oxytetracycline 10% is distributed very quickly in the blood and tissues, and very high concentrations are found during approximately 24 hours.

Indications:

Infections caused by germs susceptible to oxytetracycline such as enteritis, salmonellosis, dysentery (Vibrio), (endo)metritis, pneumonia, footrot and infections of the urinary system.

Contra-Indications:

Liver and renal insufficiencies.

Do not administer high doses to horses.

Administration/dosage:

By intramuscular or slow intravenous injection, for 3 days:

Cow : 4 ml per 100 kg b.w. daily.

Sheep, pigs : 2-4 ml per 50 kg b.w. daily.

Cats, dogs : 1 ml per 10 kg b.w. daily.

Withdrawal period:

Meat : 28 days.

Milk : 3 days.

Conservation:

At room temperature (15-25°C), protected from light.

Presentation:

Per 12 vials of 100 ml.

Per 12 vials of 50 ml

Per 6 vials of 500 ml