

Sulfadimethoxine 20%

Composition:

Each ml solution contains:
200 mg sulfadimethoxine

Action:

Sulfadimethoxine is a sulfonamide with bacteriostatic activity against Gram-positive and Gram-negative bacteria, such as Streptococci, Staphylococcus spp., E.coli, Pasteurella spp., Haemophilus spp., Klebsiella spp. and Salmonella spp. Sulfadimethoxine is rapidly absorbed from the site of application and slowly excreted.

Indications:**Cattle:**

Respiratory tract infections caused by Pasteurella spp., Salmonella dublin, Streptococcus spp., Corynebacterium pyogenes, Haemophilus somnus, Fusobacterium necrophorum; gastro-intestinal infections caused by E.coli, Salmonella sp.; genital and urinary tract infections caused by Corynebacterium spp., Streptococcus spp., E.coli, Salmonella spp.; wound infections caused by Gram-positive and Gram negative bacteria and anaerobe bacteria; mastitis caused by E.coli, Streptococcus spp., Staphylococcus spp., Klebsiella spp.; necrotic pododermatitis (foot rot) caused by Fusobacterium necrophorum.

Pigs:

Respiratory tract infections caused by Pasteurella multocida, Streptococci, Actinobacillus spp.; gastro-intestinal infections caused by E.coli, Salmonella spp. urinary and genital tract infections caused by Corynebacterium suis, Streptococci, E.coli, E. rhusiopathiae; wound infections caused by Streptococci, Corynebacterium spp., E.coli, anaerobe bacteria; mastitis caused by E.coli, Streptococci.

Administration/dosage:

By intramuscular or intravenous injection:

cattle, calves, pigs, piglets: 25 mg per kg body weight once daily, for 3 days.

Contra-Indications:

Hypersensitivity to sulfonamides; liver and kidney insufficiency.

Advised withdrawal times:

Milk : 5 days; Meat : 14 days.

Conservation:

Cool (8-15°C) and protected from light.

Presentation:

Per 12 vials of 100 ml. Per 6 vials of 250 ml