

## **Tylosin 20%**

### **Composition:**

Each ml solution contains:  
200 mg tylosin ELANCO (as tartrate)

### **Action:**

Macrolide with broad spectrum, active mainly against Gram-positive bacteria, Mycoplasma (PPLO), Actinomyces, some Spirochaetes such as Treponema hyodysenterica, Haemophilus, Erysipelotrix insidiosa, Corynebacterium diptheroides, Vibrio coli, Moraxella bovis, Listeria monocytogenes, Bacteroides, Fusobacterium.

The effective blood level is quickly obtained (1-2 hours) and lasts about 10 h. Tylosin 20% is very well distributed everywhere in the body, except maybe in the brain.

### **Indications:**

Respiratory infections, dysentery caused by Vibrio in piglets, cystitis, post-operative infections, Mycoplasma (pneumonia, CRD, arthritis, polyserositis).

### **Administration/dosage:**

By intramuscular injection:

Cattle : 3 - 5 ml per 100 kg b.w. daily.

Calves : 1.5 - 2 ml per 50 kg b.w. daily.

Pigs : 2.5 ml per 50 kg b.w. daily.

Dogs, cats : 0.5 - 1 ml per 10 kg b.w. daily for 3-5 days

### **Advised Withdrawal times:**

Meat : 8 days.

Milk : 4 days.

### **Conservation:**

Cool (8-15°C), protected from light.

### **Presentation:**

Per 12 vials of 100 ml.