

Anabolin forte

Composition:

Each ml solution contains:

50 mg nandrolone decanoate (19-nortestosterone-17-decanoate)

Action:

Stimulation of protein synthesis/prevention of protein decomposition.

Increase of volume, weight and strength of the muscles. Improvement of calcium retention and recalcification of the bones. Prolonged Action : starts from the first day and lasts till 3 to 4 weeks after injection. due to very pronounced anabolic action and very low androgenic activity Anabolin forte can be used on both male and female animals.

Indications:

Weakened animals and animals with retarded growth. Reduction of time of convalescence after chronic diseases and exhaustion. (bacterial or parasitical infections, high performances with horses and racing dogs, after exhausting delivery). Fractures, osteoporosis, deforming arthrosis, rachitis, podotrochleitis, decalcification due to high milk yield, during prolonged treatment with corticosteroids, nephrosis, uraemia etc...

Administration/dosage:

By intramuscular or subcutaneous injection:

In general: 1 mg of anabolic per kg b.w. per 3-4 weeks:

Horses, cattle : max 4 ml.

Foals, calves, pigs : max 2 ml.

Sheep, goats : max 1 ml.

Cats, dogs : max ½ ml.

Conservation:

At room temperature (15-25°C), protected from light.

Presentation:

Per 12 vials of 50 ml.

Per 25 vials of 10 ml.