

Dexpyranal

Composition:

Each ml solution contains:

140 mg isopyrin

70 mg phenylbutazone sodium

0.5 mg dexamethasone disodium phosphate.

Action and uses:

Phenylbutazone is a very active analgesic, antipyretic and antiphlogistic. Isopyrin has the same properties and increases the action of phenylbutazone. At the same time it decreases the toxicity of phenylbutazone and prolongs the activity. In this balanced composition dexpyranal is very well tolerated, even during prolonged applications. Dexamethasone is a very powerful corticosteroid with anti-inflammatory, anti-shock, anti-stress and glycogenic activities.

Indications:

All conditions that are accompanied by pain or inflammation, especially when these symptoms are due to trauma or accompanied by oedema. (Peri-)arthritis, bursitis, tendinitis, myositis, neuritis, fractures, tenosynovitis and, in combination with antibiotics, mastitis, pneumonia, Puerperal Fever.

Administration/dosage:

By intramuscular, subcutaneous or slow intravenous injection:

Horses : 20 - 40 ml iv slowly.

Cattle : 20 - 30 ml im/s.c.

Pigs : 5 - 12 ml im/s.c.

Calves : 4 - 10 ml im/s.c.

Piglets : 0.5 -1.5 ml im/s.c.

Dogs : 0.5 - 5 ml im/s.c.

Cats : 0.2 - 0.5 ml im or iv slowly

Do not administer more than 10 ml at one site in cattle (or 2 ml in dogs). Too high a dose administered at one site may cause symptoms of intolerance.

Contra-Indications:

Severe impairment of the circulatory, renal or hepatic systems. Very young animals. Peptic or intestinal ulcer, Diabetes mellitus, osteoporosis. Final stage of pregnancy.

Advised withdrawal times:

Meat : 12 days.

Milk : 4 days.

Conservation:

Cool (8-15°C), protected from light.

Presentation:

Per 12 vials of 100 ml.