

## **Erythromycin 20%**

**Composition:**

Each ml solution contains:  
200 mg erythromycin (as thiocyanate)

**Action:**

Erythromycin is a macrolide active against a wide variety of pathogenic organisms, primarily Gram-positive cocci like staphylococci, pneumococci, streptococci and enterococci. In addition it is active against other types of micro-organisms, such as Corynebacteriaceae, Haemophilus spp., Pasteurella spp. and Mycoplasma spp.

**Indications:**

Cattle, sheep: respiratory tract infections, metritis, mastitis;  
pigs: respiratory tract infections, and as an aid in the management of mastitis and metritis in sows;  
dogs and cats: respiratory tract infections and wound infections.

**Administration/dosage:**

By deep intramuscular injection;  
Cattle, sheep : 2- 4 mg per kg body weight;  
Pigs : 4-10 mg per kg body weight;  
treatment may be repeated at 24 hourly intervals if necessary.  
dogs and cats : 5-10 mg per kg body weight two times daily, during 3-5 days.

**Warnings:**

Swelling or soreness encountered at the site of injection is usually mild, transient and disappears in one or two days.

**Advised withdrawal times:**

meat: 7 days.  
milk: 3 days.

**Conservation:**

Cool (8-15°C) and protected from light.

**Presentation:**

Per 12 vials of 100 ml.