

## **Ketatroxyl**

### **Composition:**

Each ml solution contains:  
60 mg ketamine (as HCl);  
2 mg xylazine (as HCl);  
0.4 mg atropine (as sulphate).

### **Action:**

The combination of ketamine, xylazine and atropine is used to induce anaesthesia in the cat by parenteral administration. Ketamine is a quick acting general anaesthetic.

Xylazine is a potent alfa-2-adrenergic agonist and produces skeletal muscle relaxation.

The combination with ketamine prolongs duration of analgesia, reduces the dose of ketamine required and shortens recovery.

Atropine prevents salivation and other autonomic nervous effects.

### **Indications:**

Cat:

Surgery such as ovariectomy, castration, caesarian section, cheek operations, tooth extractions, eye-, nose- and ear operations, tooth cleaning, opening of abscesses.

### **Administration/dosage:**

By intramuscular and subcutaneous injection.

The recommended dose is 0.25 ml per kg body-weight (15 mg ketamine, 0.5 mg xylazine and 0.1 mg atropine per kg).

### **Contra-Indications:**

Decompensatio cordis; liver- or kidney disorders; glaucoma; head injuries.

### **Conservation:**

At room temperature (15-25°C), protected from light.

### **Presentation:**

Per 25 vials of 10 ml.