

Alfamycine 5%

Composition:

Each ml solution contains:

50 mg gentamycin base (as sulphate)

Action:

Broad spectrum antibiotic, active against most Gram-negative and certain Gram-positive germs. The action is comparable to other aminoglycosides, such as streptomycin, neomycin or kanamycin, but its sensitivity is larger.

Indications:

Infections by germs sensitive to gentamycin, such as Salmonella, Corynebacterium; colibacillosis, mastitis caused by E. coli, septicaemia, infections of the digestive and urinary tract.

Administration/dosage:

By intramuscular injection:

Horse, cattle : 8.0 ml/100 kg b.w. daily for 3-5 days.

Calf, foal, pig : 4.0 ml/ 50 kg b.w. daily for 3-5 days.

Piglet : 0.4 ml/ 5 kg b.w. daily for 3-5 days.

dog, cat : 0.4 ml/ 5 kg b.w. daily for 3-5 days.

In case of severe or chronic infection, the treatment may be repeated once or twice (half a dose every 12 h).

Contra-Indications:

Animals with kidney disorder.

Advised withdrawal times:

meat : 7 days.

milk : 3 days.

Conservation:

At room temperature (15 - 25 °C), protected from light.

Presentation:

Per 12 vials of 100 ml.