

Alfamisole 10%

Composition:

Each ml solution contains:
100 mg levamisole base (as HCl).

Action:

Levamisole demonstrates a broad spectrum activity in the removal of gastrointestinal and respiratory Nematodes. Levamisole has a paralyzing effect, with a rapid expulsion of living worms including the migrating larvae of *Ascaris suum* and *Strongyloides*. Worms expelled later are decomposed and may not be apparent in feces.

Indications:

Infections caused by gastrointestinal and respiratory nematodes susceptible to levamisole.

Pigs: *Ascaris suum* and larva migrans, *Strongyloides* spp., *Metastrongylus elongatus*, *Oesophagostomum dentatum*, *Trichuris suis* and mature *Stephanurus*, *Hyostrongylus rubidus*.

Cattle: *Dictyocaulus* spp., *Protostrongylus* spp., *Ascarides*, *Nematodirus* spp., *Strongyloides*, *Bunostomum* spp., *Oesophagostomum* spp., *Chabertina ovina*, *Strongyloides* spp.

Administration/dosage:

By a single intramuscular injection:
Cattle, sheep, goats, pigs: 1 ml per 20 kg body weight.

Advised withdrawal times:

Milk : 48 hrs.

Meat : 3 days.

Conservation:

Cool (8-15°C), protected from light.

Presentation:

Per 12 vials of 100 ml.

6 vials of 250 ml

6 vials of 500 ml